



Drug Interactions



Scott R. Penzak, Pharm.D.

Pharmacokineticist

Clinical Pharmacokinetics Research Laboratory

Clinical Center Pharmacy Department

National Institutes of Health

Drug Interactions

** Drug Interaction:*

- *The pharmacologic or clinical response to the administration of a drug combination different from that anticipated from the known effects of the two agents when given alone*
- May be harmful: toxicity, reduced efficacy
- May be beneficial: synergistic combinations, pharmacokinetic boosting, increased convenience, reduced toxicity, cost reduction

Beneficial Drug Interactions

*** Saquinavir & ritonavir**


- Saquinavir poorly absorbed, TID dosing, high pill burden (18 caps per day!)
- Combination with ritonavir results in 20-fold increase in C_{pss}
- Allows for BID dosing and decreased dose from 1200 mg TID to 1000 mg BID (1600 QD dosing is also possible)

*** Indinavir, amprenavir, atazanavir + ritonavir**

*** Cyclosporin and ketoconazole**

- Difficult to determine doses due to large interpatient variability in CYP3A (and P-gp) activity

PK Interactions between PIs: Pharmacoenhancement



Slide Courtesy of Dr. David Back. 9th Conference on Retroviruses and Opportunistic Infections. February 24-28, 2002.

Beneficial Drug Interactions: RTV + SQV




Figure 3 Plasma concentration–time profiles for subjects taking SQV–SGC monotherapy (800 mg, regimen A, ■), or 400:800 mg (regimen G, ◆) combination RTV : SQV–SGC therapy twice daily.

Epidemiology of Drug-Drug Interactions

- * True incidence difficult to determine**

- Data for drug-related hospital admissions do not separate out drug interactions, focus on ADRs

- * Most data are in the form of case reports**

- Missing or incomplete information

- * Patients receiving polypharmacy are at risk**

- 77% of HIV patients on protease inhibitors experience drug interactions

- * Difficulty in assessing role of OTC and herbal drugs in drug interactions**

- Questions regarding “active” ingredient in herbal meds

Types of Drug Interactions


*** Pharmacodynamic**

- Related to the drug's effects in the body
- One drug modulates the pharmacologic effect of another: additive, synergistic, or antagonistic

*** Pharmacokinetic**

- What the body does with the drug
- One drug alters the concentration of another
- *Usually* mediated by cytochrome P450 (CYP)

Pharmacokinetic Interactions



Drug Interactions

Absorption: food, chelation, G.I. motility, pH

Distribution: transport, protein binding

Metabolism: Phase I (CYP450), Phase II
(conjugation)


Elimination: Renal (glomerular filtration);
transport

Alterations in Absorption


*** Administration with food**

- Decreased rate of absorption; not extent (\leftrightarrow AUC):
 - * Common for many drugs; take without regard to meals
- Decreased extent of absorption (\downarrow AUC):
 - * Indinavir AUC decreased by 77% with high calorie meal; take on an empty stomach
- Increased extent of absorption (\uparrow AUC):
 - * Itraconazole (capsules) AUC increased by 66% with standard meal

Alterations in Absorption: Food Effects



Pharm Res. 1999 May;16(5):718-24.




Antimicrob Agents Chemother.
1993 Apr;37(4):778-84.

Alterations in Absorption: Chelation

* Chelation


- Irreversible binding of drugs in the GI tract
- Tetracyclines, quinolone antibiotics - ferrous sulfate (Fe^{+2}), antacids (Al^{+3} , Ca^{+2} , Mg^{+2}), dairy products (Ca^{+2})
- Usually separating administration of chelating drugs by 2+ hours decreases interaction effect

Trovafloxacin +/- Maalox®




Alterations in Absorption: GI Motility


- * **↑ GI motility:**
cisapride,
metoclopramide
- * **↓ motility:** narcotics,
antidiarrheals, high
calorie meal /
viscosity (delayed
gastric emptying)



Alterations in Absorption



Drug Interactions: Transport Proteins




Alterations in Absorption: Drug Transport


*** Efflux proteins**

- P-glycoprotein, MRP1, MRP2, OAT3
- Extrude drug from gut back into lumen limiting drug absorption
- Transporter induction may result in ↓ absorption
- Transporter inhibition may result in ↑ absorption
- Effects often difficult to assess (vs. metabolism; vs. anatomic site)
- Inhibition may be of clinical significance for drugs that are large molecules, have low bioavailability, dissolve slowly and/or incompletely (clinical significance may be overstated in medical literature)


Simplified Example of P-gp Function




LYMPHOCYTE




BRAIN




G.I. TRACT



KIDNEY




PLACENTA



LIVER

Drug Interactions: Transport Proteins

CONCEPT USING THE KIDNEY



CLINICAL APPLICATION: HEALTHY HUMAN VOLUNTEERS


	Digoxin + Placebo	Digoxin + Clarithromycin	P
AUC 0-24 hr	14	23	< .05
Cl _R (mL/min)	57	34	< .05

Alterations in Absorption: Drug Transport

*** Uptake proteins**

- OATP: located on the luminal border of enterocytes
- Transport drug across lumen and promote absorption
- Transporter inhibition may result in ↓ absorption and reduced bioavailability
- OATP substrates
 - * Pravastatin, digoxin, fexofenadine, benzylpenicillin
- OATP inhibitors
 - * Fruit juices, ritonavir, saquinavir, lovastatin, others?
- In the intestine, OATP functions *OPPOSITE* of P-gp (i.e. P-gp inhibition *INCREASES* drug absorption while OATP *DECREASES* drug absorption for compounds that are substrates of both proteins)

OATP Function: INTESTINE



Alterations in Absorption: Drug Transport




Fig 3. Mean plasma fexofenadine concentration–time profiles for persons ($n = 10$) orally administered fexofenadine (120 mg) with 300 ml water, grapefruit juice at 25% of regular strength (25% GFJ), grapefruit juice (GFJ), orange juice (OJ), or apple juice (AJ) followed by 150 ml of the same fluid every 0.5 to 3 hours (total volume, 1.2 L).

Alterations in Absorption: anion exchange resins and Gut Flora Modulation

*** Anion exchange resins (i.e. cholestyramine)**

- Form insoluble complexes with various drugs reducing their absorption
 - * Warfarin, digoxin, β -blockers, NSAIDS, others?
- Stagger dose of exchange resin with other meds
 - * Difficult due to multiple daily dosing of cholestyramine

*** Inhibition of drug-metabolizing enteric bacteria**

- Antibiotics
 - * Digoxin (*Eubacterium lentum*)
 - * Oral contraceptives (bacteria hydrolyze steroid conjugates)
 - Reports of unplanned pregnancy: causal relationship with antimicrobial administration is controversial

Alterations in Distribution: Protein Binding

“...the overall clinical importance of plasma protein binding displacement interactions continues to be overstated...”


“Despite the theoretical and experimental data to the contrary, the concept that plasma protein binding displacement is a common cause of clinically significant interactions may still be widely taught in some medical schools, often appears in textbooks and is accepted by many in the medical community and by drug regulators.”

Sansom LN & Evans AM. Drug Safety 1995;12:227-233.

Rolan PE. Br J Clin Pharmacol 1994;37:125-128.

Distribution: Drug Transport (P-gp)

^{14}C Nelfinavir +/- LY-335979 in *MDR1a* wild type Mice



● Tissue ^{14}C NFV conc. in brain
 ^{14}C NFV + LY-335979 (P-gp inhibitor)

○ Plasma ^{14}C NFV concentration
 ^{14}C NFV + LY-335979 (P-gp inhibitor)

△ Plasma ^{14}C NFV concentration
 ^{14}C NFV + vehicle

▲ Tissue ^{14}C NFV conc. in brain
 ^{14}C NFV + vehicle




Choo EF et al. Drug Metab Dispos 2000;28:655-660 Choo
EF et al. Drug Metab Dispos 2000;28:655-660.

Drug Metabolism Interactions

*** Drug metabolism**

- Chemical modification of a xenobiotic
- Phase I (functionalization RX)
 - * **Cytochrome P450 (CYP): i.e. CYP3A4, CYP2D6, CYP1A2 etc.**
- Phase II (synthetic RX)
 - * **Conjugation: i.e. glucuronidation (UGT1A1 etc.)**
- Purpose: detoxification of foreign compounds
- Anatomic sites: Liver*, Gut*, kidney, lung, brain etc.

Drug Metabolism Interactions




CYP 450 Substrates

- * **Drugs may be metabolized by a single isoenzyme**
 - Desipramine/CYP2D6; indinavir/3A4; midazolam/3A, caffeine/CYP1A2; omeprazole/CYP2C19
- * **Drugs may be metabolized by multiple isoenzymes**
 - Most drugs metabolized by more than one isozyme
 - * Imipramine: CYP2D6, CYP1A2, CYP3A4, CYP2C19
 - If co-administered with CYP450 inhibitor, some isozymes may “pick up slack” for inhibited isozyme
- * **Extensive listing + references:**
 - <http://medicine.iupui.edu/flockhart/table.htm>

CYP 450 Enzyme Inhibition

- * **Usually by competitive binding to enzyme site**
- * **Onset and offset dependent on the half-life and time to steady-state of the inhibitor**
 - Fluoxetine & CYP2D6; ritonavir and CYP3A4
- * **Time to maximum interaction effect dependent on time required for substrate drug to reach new steady-state**
- * **Mechanism-based enzyme inactivation**
 - Grapefruit juice and intestinal CYP3A content

Enzyme Inhibition



CYP 450 Inhibitors


*** The “usual suspects”**

- Cimetidine (various)
- Erythromycin, clarithromycin (3A4)
- Ketoconazole, itraconazole (3A4)
- HIV protease inhibitors (esp. ritonavir)
- Fluoxetine, paroxetine (CYP2D6)
- Nefazodone (CYP3A4)
- Grapefruit Juice (intestinal CYP3A4 only)

*** Extensive listing with references:**

- <http://medicine.iupui.edu/flockhart/table.htm>

Sildenafil (Viagra®) + Grapefruit Juice



CYP450 Inhibition

*** Key questions:**

- What is the toxic potential and therapeutic index of the substrate
 - * Terfenadine or digoxin vs sertraline**
- What are the other pathways involved in the metabolism of the substrate
 - * Zolpidem vs triazolam**
- Does the substrate have active metabolites?
 - * Codeine morphine**

CYP450 Induction

*** The “usual suspects”**

- Rifampin
- Rifabutin
- Carbamazepine
- Phenobarbital
- Phenytoin
- Nevirapine, efavirenz
- St. John's wort
- Troglitazone, pioglitazone

CYP450 Induction

- * **Gradual onset and offset**

- (involves increased DNA transcription and synthesis of new CYP enzymes)

- * **Onset and offset**

- Depends on $T_{1/2}$ of inducer, time to make new CYP proteins, and rate of degradation of CYP proteins

- * **Results in reduction of plasma concentration of substrate drugs**

- Risk of therapeutic failure
- Removal of inducer may lead to toxic concentrations of substrate
- Induction may lead to formation of toxic metabolite

CYP450 Induction

*** Mechanisms**

- **Receptor activation**

- * **Aryl hydrocarbon**


- * **Peroxisome proliferators**

- * **Constitutive androstane receptor (CAR)**



- * **Pregnane X receptor (PXR) -3A4**

- **PXR binding and activation assays: can be used to predict CYP3A4 induction**

Enzyme Induction




Induction: Influence of Ritonavir on Olanzapine Disposition in Healthy Volunteers



St. John's wort: CYP3A4 Induction Effects

● Indinavir ● Indinavir + SJW



- * 8 normal volunteers
- * Indinavir AUC determined before and after 14 days SJW 300 mg TID
- * Indinavir AUC decreased by $57 \pm 19\%$ in presence of SJW

Piscitelli SC et al. Lancet 2000;355:547-8

Predicting Drug Interactions: in vitro Screening


- * Drug development: predicting *in vivo* drug interactions from *in vitro* data Microsomes, hepatocytes, liver slices, purified CYP enzymes etc.
 - Limitations and caveats
 - * Most systems can only assess inhibition (not induction)
 - Methadone + ritonavir: discordant *in vivo* / *in vitro* results
 - * Hard to extrapolate data when drugs have multiple CYP pathways
 - * *In vitro* concentrations used may be excessively high
 - Ritonavir inhibition of MRP2

Predicting Drug Interactions: using CYP phenotypes

*** Probe + putative inhibitor or inducer**

- Measure probe (+/- metabolite(s) concentration(s)
- Ratios of metabolite:parent compound
- Examples of CYP probes
 - * **CYP1A2**: caffeine
 - * **CYP2C9**: tolbutamide; warfarin (+ vitamin K!)
 - * **CYP2C19**: S-mephenytoin; omeprazole
 - * **CYP2E1**: chlorzoxazone
 - * **CYP2D6**: dextromethorphan; debrisoquine; sparteine
 - * **CYP3A4/5**: midazolam
 - * **CYP3A4**: erythromycin


Drug Interaction Studies by Medical Division 1992-1997



PRESCRIBING FOR CARDIAC SAFETY

QTc Prolongation and Increased Cardiac Risks

Recently, the FDA and members of the medical community have raised concerns about medications associated with QTc prolongation. In normal cardiac conduction, the QTc interval lasts approximately 420 milliseconds. However, if repolarization is delayed, prolongation of the QT interval results. In general, a QTc longer than 450 milliseconds is of potential concern. Prolongation longer than 500 milliseconds indicates an elevated risk of progression to tachyarrhythmias (torsade de pointes), which may be associated with symptoms such as palpitations, dizziness, lightheadedness, and syncope; and progression to ventricular fibrillation, which can potentially cause sudden death.^{1,2} Therefore, it is important for clinicians to understand and consider this important safety issue when treating their patients.



Herb-Drug Interactions Limitations

- * **Since not regulated by FDA, safety & efficacy not required**
 - Little information available regarding drug interactions
- * **Extrapolation of data to available products difficult**
 - Independent lab tests many products (<http://www.consumerlabs.com/>)
 - 6/13 SAME preparations did not pass testing
 - * **no detectable SAME noted in one product**
 - 8/17 valerian preparations did not pass testing
 - * **4 - no detectable levels of valerenic acid**
 - * **4 - 1/2 the amount claimed on the label**

Evaluation of Specific Drug Interactions

- * **What is the time-course of the interaction**
 - Immediately or over a period of time
- * **Is it a drug class effect**
 - omeprazole vs. lansoprazole
- * **Is the interaction clinically significant**
 - Therapeutic index of drugs
 - * **Narrow or wide?**
- * **How should the interaction be managed?**
 - DC drug? Switch to another drug? Change dose?

Drug Interactions: General Tools for Evaluation and Management

- * **Familiarity with metabolic pathways**
- * **Know where to locate information on interactions**
- * **Obtain thorough medication HX at each visit**
- * **Maintain high index of suspicion when:**
 - Therapeutic response is less than expected
 - Toxic effects are present
- * **Choose drugs that are less likely to interact**
- * **Consider TDM in certain situations (anti-TB TX)**
 - Anti-TB and anti-HIV therapy

Drug Interactions: Resources

THE
END

**TABLE 5. WEB SITES WITH INFORMATION
ABOUT DRUG INTERACTIONS.**

www.dml.georgetown.edu/depts/pharmacology (Department of Pharmacology, Georgetown University Medical Center)

www.foodmedinteractions.com (food and drug interactions)

www.hivatis.org (HIV/AIDS Treatment Information Service)

www.hivdent.org (dental information)

hivinsite.ucsf.edu

www.hiv.net (in German)

www.hopkins-aids.edu (Johns Hopkins AIDS Service)

www.iapac.org (International Association of Physicians in AIDS Care)

www.hiv-druginteractions.org (Liverpool HIV Pharmacology Group)

<http://hivinsite.ucsf.edu/arvdb?page=ar-00-02&post=7>

<http://www.naturaldatabase.com>

THE
END